

CS 442

Introduction to Cryptography

Lecture 9: Example Problems on PRGs

Instructor: Aarushi Goel
Spring 2026


Pseudorandom Generators (PRG)

Definition: A deterministic algorithm G is called a pseudorandom generator if:

- * G can be computed in polynomial time.
- * $|G(x)| > |x|$
- * For every PPT adversary, $\Pr[b = b'] = \frac{1}{2} + \text{negl}(|x|)$ in the following game



Adv



Ch

$b \xleftarrow{\$} \{0,1\}$

if $b = 0$: $r \xleftarrow{\$} \{0,1\}^{\ell(n)}$

if $b = 1$: $s \xleftarrow{\$} \{0,1\}^n$
 $r = G(s)$

Example #1

Q Let $G_1: \{0,1\}^n \rightarrow \{0,1\}^{2n}$ and $G_2: \{0,1\}^n \rightarrow \{0,1\}^{2n}$ be PRGs.

Is $G(s) = G_1(s) \parallel G_2(s)$ also a PRG?

A No!

Counter example: What if G_1 and G_2 are the same PRG.

$\Rightarrow G_1(s) = G_2(s)$, the first $2n$ -bits of the output of $G(s)$ are identical to the last $2n$ -bits of the output of $G(s)$.

- This is highly unlikely to happen in a randomly sampled string.
- More formally,

$$\Pr [s_1 = s_2 \mid s \xleftarrow{\$} \{0,1\}^n, s_1 \parallel s_2 = G_1(s) \parallel G_2(s)] = 1$$

$$\Pr [s_1 = s_2 \mid s_1 \parallel s_2 \xleftarrow{\$} \{0,1\}^{4n}] = \frac{1}{2^{2n}}$$

- Here the output of G and a random string are trivially distinguishable.

Example #2

Q Let $G_1: \{0,1\}^n \rightarrow \{0,1\}^{2n}$ and $G_2: \{0,1\}^n \rightarrow \{0,1\}^{2n}$ be distinct PRGs.

Is $G(s) = \begin{cases} G_1(s) & \text{if } s \text{ is odd} \\ G_2(s) & \text{if } s \text{ is even} \end{cases}$ also a PRG?

A No!

Counter example: Let $F: \{0,1\}^{n-1} \rightarrow \{0,1\}^{2n-1}$ be a PRG. We define

We define $G_1(s_1, \dots, s_n) = F(s_1, \dots, s_{n-1}) \| s_n$ and $G_2(s_1, \dots, s_n) = F(s_1, \dots, s_{n-1}) \| \bar{s}_n$

Now, by definition of G_1 , if s is odd, last bit of $G_1(s)$ is 1.

If s is even, last bit of $G_2(s)$ is $\bar{0} = 1$.

⇒ The last bit of $G(s)$ is always 1. But this only happens with probability $\frac{1}{2}$ in a randomly sampled string.


Complement



Distinguisher / Adversary:



Checks if the last
bit of r is 1.
If so, set $b' = 1$
else, set $b' = 0$



$b \leftarrow \{0,1\}$
if $b = 0$: $r \leftarrow \{0,1\}^{2n}$
if $b = 1$: $s \leftarrow \{0,1\}^n$
 $r = G(s)$

$$\begin{aligned} \Pr[b=b'] &= \Pr[b=b'=0] + \Pr[b=b'=1] \\ &= \frac{1}{2} \times \Pr[b'=0|b=0] + \frac{1}{2} \times \Pr[b'=1|b=1] \\ &= \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times 1 = \frac{1}{2} + \frac{1}{4} \end{aligned}$$

→ non-negligible advantage.

Example #3

Q Let $G_1: \{0,1\}^n \rightarrow \{0,1\}^{2n}$ and $G_2: \{0,1\}^{2n} \rightarrow \{0,1\}^{4n}$ be PRGs. Prove that the following function is also a PRG: $F: \{0,1\}^n \rightarrow \{0,1\}^{4n}$, $F(x) = G_2(G_1(x))$

A. We need to show that the following two distributions are computationally indistinguishable:

$$\{ G_2(G_1(x)) ; x \xleftarrow{\$} \{0,1\}^n \}$$

$$\{ s \xleftarrow{\$} \{0,1\}^{4n} \}$$

Consider the following hybrids:

$$H_1: \left\{ G_2(G_1(x)) ; x \xleftarrow{\$} \{0,1\}^n \right\}$$

$$H_2: \left\{ G_2(r) ; r \xleftarrow{\$} \{0,1\}^{2n} \right\}$$

$$H_3: \left\{ s \xleftarrow{\$} \{0,1\}^{4n} \right\}$$

Following the hybrid lemma, it suffices for us to show that

$$H_1 \approx_c H_2 \quad \text{and} \quad H_2 \approx_c H_3.$$

* $H_2 \approx_c H_3$ follows directly from pseudorandomness of G_2 .

* Let us focus on proving $H_1 \approx_c H_2$ using a proof by reduction.

Proof by Reduction

$$H_1: \{ G_2(G_1(x)) ; x \xleftarrow{\$} \{0,1\}^n \} \quad H_2: \{ G_2(r) ; r \xleftarrow{\$} \{0,1\}^{2n} \}$$

* To prove that $H_1 \approx_c H_2$, we will use the following line of reasoning:

1. Let us assume for the sake of contradiction that \exists a non-uniform PPT adversary A , can distinguish between H_1 and H_2 with some non-negligible probability.
2. We will use A to construct another non-uniform PPT adversary B who can break pseudorandomness of G_1 with non-negligible advantage.
3. But we know that G_1 is a PRG. Therefore no such adversary B can exist. Hence we arrive at a contradiction implying that our assumption was incorrect.


Proof by Reduction: How to construct B using A?

$$H_1: \{ G_1(x) ; x \xleftarrow{\$} \{0,1\}^n \} \quad H_2: \{ G_2(r) ; r \xleftarrow{\$} \{0,1\}^{2n} \}$$

Recall the game-based definition of PRG.



Adv



Ch

$$b \xleftarrow{\$} \{0,1\}$$


if $b=0$: $r \xleftarrow{\$} \{0,1\}^{\ell(n)}$
if $b=1$: $s \xleftarrow{\$} \{0,1\}^n$
 $r = G(s)$

$$\Pr [b = b'] = \frac{1}{2} + \text{negl}(\lambda)$$

To prove: $H_1 \approx_c H_2$

$$H_1: \left\{ G_2(G_1(x)) ; x \xleftarrow{\$} \{0,1\}^n \right\}$$

$$H_2: \left\{ G_2(r) ; r \xleftarrow{\$} \{0,1\}^{2n} \right\}$$




If s is pseudorandom, then input to A is distributed identically to a sample from H_1 , else it is identically distributed to a sample from H_2 .

To prove: $H_1 \approx_c H_2$

$$H_2: \left\{ G_2(G_1(x)) ; x \xleftarrow{\$} \{0,1\}^n \right\}$$

$$H: \left\{ G_2(r) ; r \xleftarrow{\$} \{0,1\}^{2n} \right\}$$



\Rightarrow If **A** succeeds with non-negligible advantage $\mu(n)$, then **B** also succeeds with the same non-negligible advantage $\mu(n)$.
This is a contradiction!

Proofs by Reduction: Key Points

- * Here are 4 important things that must keep in mind for a valid reduction:
1. **Input Mapping:** How to map the input that the outer adversary **B** receives from the challenger to an input for the inner adversary **A**.
 2. **Input Distribution:** Does the above input mapping provide the right distribution of inputs that **A** expects.
 3. **Output Mapping:** How do we map the output that **A** provides to an output for **B**.

4. **Win Probability**: When we assume existence of A , we also assume that A wins with some non-negligible advantage $\mu(n)$. What is the probability or advantage with which B wins in terms of $\mu(n)$, given the above input/output mappings?